



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : A61K 7/08, 7/50, 7/02 C11D 1/94	A1	(11) International Publication Number: WO 92/13513 (43) International Publication Date: 20 August 1992 (20.08.92)
(21) International Application Number: PCT/US92/00749 (22) International Filing Date: 30 January 1992 (30.01.92) (30) Priority data: 654,177 12 February 1991 (12.02.91) US (71) Applicant: RICHARDSON-VICKS INC. [US/US]; One Far Mill Crossing, Shelton, CT 06484 (US). (72) Inventors: LINARES, Carlos, Gabriel ; 20 Brownley Drive, Stamford, CT 06905 (US). DECKNER, George, Endel ; 22 Powder Mill Lane, Trumbull, CT 06611 (US). ST. JOHN, Lucie, Anita ; 171 Sunnyside Avenue, Waterbury, CT 06708 (US).		(74) Agent: REED, T., David; The Procter & Gamble Company, Ivorydale Technical Ctr., 5299 Spring Grove Avenue, Cincinnati, OH 45217-1087 (US). (81) Designated States: AT, AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CI (OAPI patent), CM (OAPI patent), CS, DE, DE (European patent), DK, DK (European patent), ES, ES (European patent), FI, FR (European patent), GA (OAPI patent), GB, GB (European patent), GN (OAPI patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (European patent), MC (European patent), MG, ML (OAPI patent), MN, MR (OAPI patent), MW, NL, NL (European patent), NO, PL, RO, RU, SD, SE, SE (European patent), SN (OAPI patent), TD (OAPI patent), TG (OAPI patent). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: COSMETIC COMPOSITIONS (57) Abstract An aqueous foaming cosmetic composition comprising: (a) from 0.1 to 7 % of an imidazolinium amphoteric surfactant, (b) from about 0.1 % to about 5.0 % of a polyol alkoxy ester wherein the ratio of (a):(b) is from about 15:1 to about 1:1 and wherein the cleansing composition has a viscosity of at least about 150 cps (Brookfield RVT, Spindle No. TB, 10 rpm, 25 °C). The composition has improved foam stability, together with excellent cleansing performance and mildness. It is suitable for use as make-up and facial cleansers, foam bath, shower products, shampoos etc.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	ML	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

COSMETIC COMPOSITIONSTECHNICAL FIELD

5 The present invention relates to improved cosmetic cleansing compositions, and in particular to foaming cosmetic compositions suitable for cleansing the skin and/or the hair and which may be used, for example, in the form of make-up removal and facial cleansers, bath
10 foams, shower products, shampoos and the like.

BACKGROUND OF THE INVENTION

 Foaming cosmetic compositions must satisfy a number of criteria including cleansing power, foaming properties and mildness/low irritancy with respect to the skin, hair and the ocular mucosae.

15 Skin is made up of several layers of cells which coat and protect the keratin and collagen fibrous proteins that form the skeleton of its structure. The outermost of these layers, referred to as the stratum corneum, is known to be composed of 250 Å diameter protein bundles surrounded by 80 Å thick bilayers of epidermal lipids and
20 water. Anionic surfactants can penetrate the stratum corneum membrane and, by delipidization (i.e. removal of the lipids from the stratum corneum), destroy its integrity. This destruction of the stratum corneum bilayers can lead to dry rough skin and may eventually permit the surfactant to interact with the viable epidermis, creating irritation.
25

 Ideal cosmetic cleansers should cleanse the skin or hair gently, causing little or no irritation without defatting and or drying the skin and without leaving skin taut after frequent use. Most lathering soaps, liquids and bars fail in this respect.

30 Certain synthetic surfactants are known to be mild. However, a major drawback of most mild synthetic surfactant systems, when formulated for skin cleansing, is poor lather performance compared to the highest bar soap standards (bars which are rich in coconut soap and superfatted). On the other hand, the use of known high sudsing
35 anionic surfactants with lather boosters can yield acceptable lather volume and quality. Unfortunately, however, the highest sudsing anionic surfactants are, in fact, poor in clinical skin mildness. Surfactants that are among the mildest, such as sodium lauryl glyceryl ether sulfonate, (AGS), are marginal in lather. These two facts make

-2-

the balancing of the surfactant selection and the lather and skin feel benefit a delicate process.

5 Rather stringent requirements for cosmetic cleansers limit the choice of surface-active agents, and final formulations represent some degree of compromise. Mildness is often obtained at the expense of effective cleansing, or lathering may be sacrificed for either mildness, product stability, or both.

10 Thus a need exists for foaming cosmetic compositions which will produce a foam which is abundant, stable and of high quality (compactness), which are effective skin and hair cleansers and which are very mild to the skin, hair and ocular mucosae.

15 It has been found that the use of specific polyol alkoxy esters in combination with specific amphoteric surfactants provide cosmetic cleansing compositions with significantly improved dermal mildness benefits and also good physical characteristics such as foaming.


It is therefore an object of the present invention to provide an improved cosmetic cleansing composition which thoroughly cleanses the skin and hair and which is very mild to the skin, hair and ocular mucosae.

20 It is a further object of the present invention to provide an improved cosmetic cleansing composition which is mild and which will produce a foam which is abundant, stable and of high quality, and which effectively cleans skin and hair.

SUMMARY OF THE INVENTION

25 The present invention relates to foaming cosmetic compositions suitable for cleansing the skin or hair and which may be used as make-up removers and facial cleansers, bath foams, shower products, shampoos, and the like comprising:

30 (a) from about 0.1% to about 7.0% by weight on a solids basis of a first amphoteric surfactant of formula I selected from imidazolinium derivatives:



-3-

wherein R_1 is C_8 - C_{22} alkyl or alkenyl, R_2 is hydrogen or CH_2COOM , Z is H, CH_2COOM , CH_2CH_2COOM , or $CH_2CHOHCH_2SO_3M$, Y is H, CH_2COOM , CH_2CH_2COOM , or $CH_2CHOHCH_2SO_3M$ and M is H, alkali metal, alkaline earth metal, ammonium or alkanol-ammonium;

- (b) from about 0.1% to about 5.0% of a polyol alkoxy ester; and
(c) from about 60% to about 99.5% water;

wherein the ratio of (a):(b) is from about 15:1 to about 1:3
and wherein the cleansing composition has a viscosity of at least
about 150 cps (Brookfield RVT, Spindle No TB, 10 rpm, 25°C).


All concentrations and ratios herein are by weight of total composition and all measurements are at 25°C, unless otherwise specified.

DETAILED DESCRIPTION OF THE INVENTION

The invention relates to a foaming cosmetic composition with superior mildness and excellent lathering characteristics (abundance, quality) and good cleansing ability. The foaming cosmetic composition take the form of a viscous liquid, paste or gel which has the advantage that it can be easily and commercially packaged in and dispensed from pump bottles or from tubes by squeezing. Gel form compositions are defined herein as those which have a viscosity (Brookfield RVT, Spindle No. TB, 10 rpm, 25°C) of at least 10,000 cps. Preferred from the viewpoint of flowability are gel compositions having a viscosity in the range from about 10,000 to about 50,000 cps, more preferably from about 10,000 to about 20,000 cps. Viscous liquids or pastes are defined herein as those having a viscosity (same conditions) of at least about 150 cps, preferably at least about 500 cps and less than about 10,000 cps, and more preferably between about 5,000 cps and about 9,000 cps.

Surfactant

The essential surfactant component of the compositions of the present invention is an amphoteric surfactant selected from imidazolinium surfactants of formula I




wherein R_1 is C₈-C₂₂ alkyl or alkenyl, R_2 is hydrogen or CH_2COOM , Z is H, CH_2COOM , CH_2CH_2COOM , or $CH_2CHOHCH_2SO_3M$, Y is H, CH_2COOM , CH_2CH_2COOM , or $CH_2CHOHCH_2SO_3M$ and M is H, alkali metal, alkaline earth metal, ammonium or alkanol-ammonium;

These amphoteric surfactants are present in the compositions of the present invention at a level of from about 0.1% to about 7%, preferably from about 0.1% to about 5%, more preferably from about 0.1% to about 3% and most preferably from about 0.1% to about 2.5%, on a solids basis.

Examples of suitable amphoteric surfactants for use herein include compounds in which R_1 is C₈H₁₇ (especially iso-capryl), C₈H₁₉, C₁₁H₂₃ and C₁₂H₂₅ alkyl. Especially preferred are the compounds in which R_1 is C₁₂H₂₅, Z is CH_2CO_2M , Y is CH_2COOM and R_2 is H. These surfactants are all fully described in Miranol Products for Cosmetics and Toiletries Technical and Product Development, Tenth Edition (1987); which is incorporated by reference herein.


It will be understood that a number of commercially-available amphoteric surfactants of this type are manufactured and sold in the form of complexes with anionic surfactants, especially those of the sulfated C₈-C₁₈ alcohol or C₈-C₁₈ acyl glyceride types. More preferably, the compositions comprise the surfactant component along with a premix or complex of the optional amphoteric surfactant and anionic surfactant in an equivalent ratio of about 1:1 in order to provide approximate electroneutrality.

In a preferred embodiment of the present invention, the compositions comprise a second amphoteric surfactant being selected from aminoalkanoates of formula II



II

iminodialkanoates of formula III



III

and mixtures thereof, wherein n and m are numbers from 1 to 4, and R_1 and M are independently selected from the groups specified in I above.

The compositions herein can also comprise other optional surfactant components, notably, anionic surfactants. It is an important feature of the invention, however, that the combined concentration of the first and the optional second amphoteric surfactants is at least about 30% by weight of the total surfactant concentration, this being important from the viewpoint of achieving optimum lathering characteristics. In preferred compositions, the mixture of the first and second amphoteric surfactants comprises at least about 60%, more preferably at least about 75% by weight of the total surfactant composition.

Examples of suitable optional amphoteric surfactants for use herein include salts, especially the triethanolammonium salts and salts of N-lauryl-beta-aminopropionic acid and N-laurylimino-dipropionic acid.

The compositions herein preferably contain from about 1% to about 10% by weight, more preferably from about 1.5% to about 5% by weight of each of the first and second amphoteric surfactants. The weight ratio of first amphoteric surfactant : second amphoteric surfactant is preferably from about 10:1 to about 1:10, more preferably from about 5:1 to about 1:5, and especially from about 3:1 to about 1:3.

A preferred optional surfactant in the compositions herein is an anionic surfactant. This is preferably present in a level of from about 0.1% to about 10%, more preferably from about 1.0% to about 8%, and especially from about 4% to about 6% by weight. Preferred anionic surfactants for inclusion herein, other than the alkyl and acylglyceride sulfates mentioned above, are the fatty acid condensation products of proteins, degraded proteins or amino acids, or mixtures of such condensation products. In highly preferred embodiments, the fatty acid condensation products are selected from:

- (i) condensation products of C₈-C₁₂, preferably C₁₀-C₁₈ fatty acids with hydrolysed proteins
- (ii) fatty acid sarconsinates derived from C₈-C₂₂, preferably C₁₀-C₁₈ fatty acids, and
- (iii) mixtures thereof.

However, the total level of surfactant in the compositions herein should generally lie in the range from about 1% to about 20% by weight, preferably from about 1% to about 15% by weight, and especially from about 2% to about 7% by weight.

Polyol Alkoxy Ester

The compositions of the present invention also essentially comprise a polyol alkoxy ester. These non-ionic compounds are high molecular weight (from about 4,000 to about 8,000 preferably from about about
5 6,000 to about 8,000) branched polyethylene glycol compounds. A highly preferred polyol alkoxy ester is available from Croda, Inc. under the trade name Crothix. The polyol alkoxy ester is present in the composition at a level of from about 0.1% to about 5%, preferably from about 0.5% to about 3%, and most preferably from about 1% to
10 about 2%.

The amphoteric surfactant and polyol alkoxy ester are present in a ratio of amphoteric surfactant:polyol alkoxy ester of from about 10:1 to about 1:3, preferably from about 5:1 to about 1:3 and most preferably from about 3:1 to about 1:3.

Optional Components

The compositions of the invention may also contain additional thickeners at a level preferably from about 0.1% to about 10%, more preferably from about 0.1% to about 5%, and especially from about 0.3% to about 4%. The thickener preferably has a viscosity (1% aqueous
20 solution, 25°C, Brookfield RVT Spindle No TB, 5 rpm) of at least about 4000 cps, more preferably at least about 10,000 cps.

Neutralizing agents suitable for use in neutralizing acidic group containing hydrophilic thickeners herein including sodium hydroxide, potassium hydroxide, ammonium hydroxide, monoethanolamine, diethanol-
25 amine and triethanolamine.

The compositions of the invention can optionally include a hair or skin moisturizer. The preferred level of moisturizer is from about 1% to about 20% by weight. In preferred embodiments, the moisturizer is nonocclusive and is selected from:

- 30 1. water-soluble liquid polyols;
2. essential amino acid compounds found naturally occurring in the stratum corneum of the skin; and
3. water-soluble nonocclusives and mixtures thereof.

Some examples of more preferred nonocclusive moisturizers are
35 glycerine, polyethylene glycol, propylene glycol, sorbitol, polyethylene glycol and propylene glycol esters of methyl glucose (e.g. methyl gluxan-20), polyethylene glycol and propylene glycol esters of lanolin alcohol (e.g. Solulan-75), sodium pyrrolidone carboxylic acid, lactic

acid, urea, L-proline, guanidine, pyrrolidone and mixtures thereof. Of the above, glycerine is highly preferred.

Examples of other water-soluble nonocclusive moisturizers include water-soluble hexadecyl, myristyl, isodecyl or isopropyl esters of adipic, lactic, oleic, stearic, isostearic, myristic or linoleic acids, as well as many of their corresponding alcohol esters (sodium isostearoyl-2-lactylate, sodium capryl lactylate), hydrolyzed protein and other collagen-derived proteins, aloe vera gel and acetamide MEA.

The compositions of the invention can additionally comprise from about 0.05% to about 5% by weight of cationic or nonionic polymeric hair or skin conditioning agents. Representative classes of polymeric hair or skin conditioning agents include cationic and nonionic polysaccharides; cationic and nonionic homopolymers and copolymers derived from acrylic and/or methacrylic acid; cationic and nonionic cellulose resins; cationic copolymers of dimethyldiallylammonium chloride and acrylic acid; cationic homopolymers of dimethyldiallylammonium chloride; cationic polyalkylene and ethoxypolyalkene imines, and mixtures thereof.

By way of exemplification, cationic polymeric conditioning agents preferred for use herein include cationic guar gums such as hydroxypropyl trimethyl ammonium guar gum (d.s. of from 0.11 to 0.22) available commercially under the trade names Jaguar C-14-S(RTM) and Jaguar C-17(RTM), and also Jaguar C-16(RTM), which contains hydroxypropyl substituents (d.s. of from 0.8 - 1.1) in addition to the above-specified cationic groups, and quaternized cellulose esters available commercially under the trade names Ucare Polymer JR and Celquat. Other suitable cationic polymers are homopolymers of dimethyldiallylammonium chloride available commercially under the trade name Merquat 100, copolymers of dimethyl aminoethylmethacrylate and acrylamide, copolymers of dimethyldiallylammonium chloride and acrylamide, available commercially under the trade names Merquat 550 and Merquat S, quaternized vinyl pyrrolidone acrylate or methacrylate copolymers of amino alcohol available commercially under the trade name Gafquat, and polyalkylenimines such as polyethylenimine and ethoxylated polyethylenimine.

A number of additional optional materials can be added to the compositions of the invention. Such materials include keratolytic agents such as salicylic acid; proteins and polypeptides and derivatives thereof; water-soluble or solubilizable preservatives such as

Germall 115, methyl, ethyl, propyl and butyl esters of hydroxybenzoic acid, EDTA, Euxyl (RTM) K400, Bronopol (2-bromo-2-nitropropane-1, 3-diol); anti-bacterials such as Irgasan (RTM), phenoxyethanol and phenoxypropanol (preferably at levels of from about 0.2% to about 5%);
 5 other moisturizing agents such as hyaluronic acid, chitin, and starch-grafted sodium polyacrylates such as Sanwet (RTM) IM-1000, IM-1500 and IM-2500 available from Celanese Superabsorbent Materials, Portsmouth, VA, USA and described in USA-A-4,076,663; colouring agents; Pearling agents; perfumes and perfume solubilizers etc. Water is also present
 10 at a level of from about 50% to about 98.9%, preferably from about 70% to about 95% by weight of the compositions herein.

The pH of the compositions is preferably from about 4 to about 9, more preferably from about 4.5 to about 7.

15 The following examples further describe and demonstrate embodiments within the scope of the present invention. The examples are given solely for the purpose of illustration and are not to be construed as limitations of the present invention, as many variations thereof are possible without departing from the spirit and scope of the invention.

Ingredients are identified by chemical or CTFA name.

20

Example I

A facial cleansing composition of the present invention is made as follows:

	<u>Ingredient</u>	<u>% w/w</u>
	<u>Phase A</u>	
25	Water	q.s.
	Polyquaternium -10	0.50
	<u>Phase B</u>	
	Potassium Coco Hydrolyzed Animal Protein ²	15.0
30	Amphoteric Surfactant I ³	5.1
	Amphoteric Surfactant II ⁴	6.8
	Amphoteric Surfactant III ⁵	7.3
	<u>Phase C</u>	
	Glycerin	3.0
35	Polyol Alkoxy Ester ⁶	1.6
	PEG-120 Methyl Glucose Dioleate ⁷	0.6
	<u>Phase D</u>	
	Preservative (phenoxyethanol)	0.4

-9-

Phase E

Water	1.0
Na ₄ EDTA	0.1

- 5 1 Available as Polymer JR 400
 2 Available as Lamepon S
 3 Cocoamphocarboxyglycinate (and) Na Lauryl Sulfate (and)
 Hexylene Glycol
 available as Miranol 2MCA MOD
 10 4 Na Lauryl Sarcosinate
 available as Hamposyl- L-30
 5 Na Lauriminodipropionate
 available as Mirataine H2C-HA
 6 Available as Crothix
 15 7 Available as Glucamate DOE-120

The water is heated to 65°C and the polyquaternium-10 is added to the water to form Phase A. The Phase B ingredients are added sequentially to this phase. Separately, the Phase C components are heated to 65°C. Phase C is combined with this mixture and then cooled to 40°C. Phase D and Phase E are added to this mix to form the resultant cosmetic composition.

- Application of approximately 2 grams of the resulting gel/viscous liquid with water is useful for topical application as a cleanser to remove, for example, dirt and oil as well as difficult to remove make-up, waterproof mascara and the like.

Example II

- A facial cleansing composition of the present invention is made by combining the following ingredients utilizing conventional mixing techniques as described above in Example I.

<u>Ingredient</u>	<u>% w/w</u>
<u>Phase A</u>	
Water	q.s.
Polyquaternium -10 ¹	0.50
<u>Phase B</u>	
Amphoteric Surfactant I ²	15.1
Amphoteric Surfactant II ³	4.0
<u>Phase C</u>	
Glycerin	3.0

-10-

	<u>Phase B</u>	
	Amphoteric Surfactant I ²	15.1
	Amphoteric Surfactant II ³	4.0
	<u>Phase C</u>	
5	Glycerin	3.0
	Polyol Alkoxy Ester ⁴	1.6
	PEG-120 Methyl Glucose Dioleate ⁵	0.6
	<u>Phase D</u>	
	Preservative (phenoxyethanol)	0.4
10	<u>Phase E</u>	
	Water	1.0
	Na ₄ EDTA	0.1
15	1 Available as Polymer JR 400	
	2 Na Lauriminodipropionate available as Mirataine H2C-HA	
	3 Cocoamphocarboxyglycinate (and) Na Lauryl Sulfate (and) Hexylene Glycol available as Miranol 2MCA MOD	
20	4 Available as Crothix	
	5 Available as Glucamate DOE-120	

Application of approximately 2 grams of the resulting gel/viscous liquid with water is useful for topical application as a cleanser to remove, for example, dirt and oil as well as difficult to remove make-up, waterproof mascara and the like.

Example III

A facial cleansing composition of the present invention is made by combining the following ingredients utilizing conventional mixing techniques as described above in Example I.

	<u>Ingredient</u>	<u>% w/w</u>
	<u>Phase A</u>	
	Water	q.s.
35	Polyquaternium -10 ¹	0.50
	<u>Phase B</u>	
	Amphoteric Surfactant I ²	15.1
	Amphoteric Surfactant II ³	4.0

-11-

	<u>Phase C</u>	
	Glycerin	3.0
	Polyol Alkoxy Ester ⁴	1.0
5	Na Acrylate/Stearate 20 Methacrylate polymer ⁵	3.0
	<u>Phase D</u>	
	Preservative	0.4
	<u>Phase E</u>	
	Water	1.0
10	Na ₄ EDTA	0.1

1 Available as Polymer JR 400

2 Na Lauriminodipropionate
available as Deriphat 160C

15 3 Cocoamphocarboxyglycinate (and) Na Lauryl Sulfate (and)
Hexylene Glycol
available as Miranol 2MCA MOD

4 Available as Crothix

5 Available as Aculyn 22

20 Application of approximately 2 grams of the resulting gel/viscous
liquid with water is useful for topical application as a cleanser to
remove, for example, dirt and oil as well as difficult to remove
make-up, waterproof mascara and the like.

25 Example IV

A facial cleansing composition of the present invention is made by
combining the following ingredients utilizing conventional mixing
techniques as described above in Example I.

30	<u>Ingredient</u>	<u>% w/w</u>
	<u>Phase A</u>	
	Water	q.s.
	Polyquaternium -10 ¹	0.50
	<u>Phase B</u>	
35	Amphoteric Surfactant I ²	8.0
	Amphoteric Surfactant II ³	8.0

-12-

Phase C

Glycerin	15.0
Polyol Alkoxy Ester ⁴	1.5
PEG-120 Methyl Glucose Dioleate ⁵	0.5

Phase D

Preservative (phenoxyethanol)	0.4
-------------------------------	-----

Phase E

Water	1.0
Na ₄ EDTA	0.1

1 Available as Polymer JR 400

2 Lauroamphodiacetate (and) Na Trideceth Sulfate
available as Miranol BT

3 Na Lauriminodipropionate

available as Mirataine H2C-HA

4 Available as Crothix

5 Available as Glucomate DOE-120


Application of approximately 2 grams of the resulting gel/viscous liquid with water is useful for topical application as a cleanser to remove, for example, dirt and oil as well as difficult to remove make-up, waterproof mascara and the like.

WHAT IS CLAIMED:

-13-

Claims:

1. A foaming cosmetic cleansing composition comprising;
 (a) from about 0.1% to about 7.0% by weight on a solids basis of an amphoteric surfactant selected from imidazolinium derivatives of formula I



wherein R₁ is C₈-C₂₂ alkyl or alkenyl, R₂ is hydrogen or CH₂COOM, Z is H, CH₂COOM, CH₂CH₂COOM, or CH₂CHOHCH₂SO₃M, Y is H, CH₂COOM, CH₂CH₂COOM, or CH₂CHOHCH₂SO₃M and M is H, alkali metal, alkaline earth metal, ammonium or alkanol-ammonium;


- (b) from about 0.1% to about 5.0% of a polyol alkoxy ester; and
 (c) from about 60% to about 99.5% water;

wherein the ratio of (a):(b) is from about 15:1 to about 1:1 and wherein the cleansing composition has a viscosity of at least about 150 cps (Brookfield RVT, Spindle No TB, 10 rpm, 25°C).

2. A cosmetic composition according to Claim 1 which further comprises from about 0.5% to about 16% by weight of a second amphoteric surfactant selected from aminoalkanoates of formula II



iminodialkanoates of formula III



and mixtures thereof, wherein R₁ is C₈-C₂₂ alkyl or alkenyl, R₂ is hydrogen or CH₂CH₂M, Z is CO₂M or CH₂CO₂M and M is H, alkali metal, alkaline earth metal, ammonium or alkanol-ammonium; and wherein n and m are numbers from 1 to 4, and

wherein the mixture of first and second amphoteric surfactants comprises at least about 30% by weight of the total surfactant component.


3. A cosmetic composition according to Claim 2 which comprises from about 0.5% to about 2.0% of the polyol alkoxy ester and wherein the total surfactant concentration is from about 1% to about 20% by weight of the cleansing composition.
4. A cosmetic composition according to Claim 3 wherein the ratio of amphoteric surfactant:polyol alkoxy ester is from about 10:1 to about 1:3 and further comprises from about 1% to about 10% by weight of each of the first and second amphoteric surfactants.
5. A cosmetic composition according to Claim 4 wherein the weight ratio of first amphoteric surfactant:second amphoteric surfactant is from about 10:1 to about 1:10.
6. A cosmetic composition according to Claim 5 wherein the weight ratio of first amphoteric surfactant:second amphoteric surfactant is from about 3:1 to about 1:3, and wherein the composition additionally comprises from about 0.1% to about 10%, by weight of anionic surfactant.
7. A cosmetic composition according to Claim 6 wherein the ratio of amphoteric surfactant:polyol alkoxy ester is from about 3:1 to about 1:3 and comprises a premix or complex of the first amphoteric surfactant and anionic surfactant in an equivalent ratio of about 1:1.
8. A cosmetic composition according to Claim 8 wherein the fatty acid condensation product is selected from
 - (i) a condensation product of C₈-C₁₂, preferably C₁₀-C₁₈ fatty acids with hydrolysed proteins,
 - (ii) fatty acid sarcosinates derived from C₈-C₂₂, preferably C₁₀-C₁₈ fatty acids, and
 - (iii) mixtures thereof.

9. A cosmetic composition according to Claim 8 having a viscosity of from about 150 to about 20,000 cps (Brookfield RVT, Spindle No. TB, 10 rpm, 25°C).
10. A cosmetic composition according to Claim 9 wherein the aqueous cleansing composition comprises from about 1% to about 20% of a hair or skin moisturizer.
11. A cosmetic composition according to Claim 10 wherein the moisturiser is nonocclusive and is selected from:
 - (i) water-soluble liquid polyols;
 - (ii) essential amino acid compounds found naturally occurring in the stratum corneum of the skin; and
 - (iii) water-soluble nonpolyol nonocclusives and mixtures thereof.
12. A cosmetic composition according to Claim 11 wherein the moisturiser is selected from glycerin, polyethylene glycol, propylene glycol, sorbitol, polyethylene glycol and propylene glycol esters of methyl glucose, polyethylene glycol and propylene glycol esters of lanolin alcohol, sodium pyrrolidone carboxylic acid, lactic acid, L-proline and mixtures thereof.
13. A cosmetic composition according to Claim 12 wherein the moisturiser is glycerin.
14. A cosmetic composition according to Claim 13 additionally comprising from about 0.05% to about 5% by weight of a cationic or nonionic polymeric hair or skin conditioning agent.
15. A cosmetic composition according to Claim 14 wherein the polymeric hair or skin conditioning agent is selected from cationic and nonionic polysaccharides; cationic and nonionic homopolymers and copolymers derived from acrylic and/or methacrylic acid; cationic and nonionic cellulose resins; cationic copolymers of dimethyldiallylammonium chloride and acrylic acid; cationic homopolymers of dimethyldiallylammonium chloride; cationic polyalkylene and ethoxypolyalkylene imines, and mixtures thereof.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 92/00749

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 A61K7/08;	A61K7/50;	A61K7/02; C11D1/94
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	A61K ; C11D	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
P,A	WO,A,9 117 237 (THE PROCTER & GAMBLE COMPANY) 14 November 1991 see the whole document	1-15
P,A	WO,A,9 111 984 (THE PROCTER & GAMBLE COMPANY) 22 August 1991 see the whole document	1-15
A	EP,A,0 250 181 (HELENE CURTIS INDUSTRIES INC.) 23 December 1987 see page 2, line 14 - page 2, line 37; claims	1
<p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
12 JUNE 1992	22.05.92	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	COUCKUYT P.J.R. 	

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

US 9200749
SA 57774

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 12/06/92

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9117237	14-11-91	AU-A- 7560091 CN-A- 1055109	27-11-91 09-10-91
WO-A-9111984	22-08-91	AU-A- 7303391	03-09-91
EP-A-0250181	23-12-87	AU-B- 596187 AU-A- 7377087 JP-A- 63022900	26-04-90 17-12-87 30-01-88

EPO FORM P0079

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82